

What to do if you have confirmed or suspected Coronavirus disease (COVID-19)

If you are sick and have been diagnosed with COVID-19 or suspected to have COVID-19 because you have been exposed to someone with COVID-19, follow the steps below to help prevent the disease from spreading to people in your home and community.

Symptoms of COVID-19

The most common symptoms of COVID-19 are fever, cough and shortness of breath. If you have been exposed to someone with laboratory confirmed COVID-19 and are experiencing fever with either cough or shortness of breath, you might have COVID-19. You can contact your doctor to see if you need to be tested. If you have tested positive for COVID-19 or are suspected to have COVID-19 but are not tested, you should follow the below instructions.

Stay home except to get medical care

You should restrict activities outside your home, except for getting medical care. Do not go to work, school, or public areas. Avoid using public transportation, ride-sharing, or taxis.

Separate yourself from other people and animals in your home

- **People:** As much as possible, you should stay in a specific room and away from other people in your home. Also, you should use a separate bathroom, if available.
- **Animals:** You should restrict contact with pets and other animals while sick. When possible, have another member of your household care for your animals while you are sick; if you must care for your pet, wash your hands before and after you interact with pets and wear a facemask, if possible. See COVID-19 and Animals for more information.

Call ahead before visiting your doctor

If you have a medical appointment, call the healthcare provider and tell them that you have or may have COVID-19. This will help the healthcare provider's office take steps to keep other people from getting infected or exposed.

Wear a facemask

You should wear a facemask, if possible, when you are around other people (e.g., sharing a room or vehicle) or pets and before you enter a healthcare provider's office. If you are not able to wear a facemask (for example, because it causes trouble breathing), then people who live with you should not be in the same room with you, or they should wear a facemask if they enter your room.

Cover your coughs and sneezes

Cover your mouth and nose with a tissue when you cough or sneeze. Throw used tissues in a lined trash can; immediately clean your hands as described below.

Clean your hands often

Wash your hands often with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Soap and water is preferred if hands are visibly dirty. Avoid touching your eyes, nose, and mouth with unwashed hands.

Avoid sharing personal household items

You should not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people or pets in your home. After using these items, they should be washed thoroughly with soap and water and dried before use by others.

What to do if you have confirmed or suspected Coronavirus disease (COVID-19)

Clean all “high-touch” surfaces every day

High touch surfaces include counters, tabletops, doorknobs, bathroom fixtures, toilets, phones, keyboards, tablets, and bedside tables. Also, clean any surfaces that may have blood, stool, or body fluids on them. Use a household cleaning spray or wipe, according to the label instructions. Labels contain instructions for safe and effective use of the cleaning product including precautions you should take when applying the product, such as wearing gloves and making sure you have good ventilation during use of the product.

Monitor your symptoms

Seek prompt medical attention if your illness is worsening (e.g., difficulty breathing). Before seeking care, call your healthcare provider and tell them that you have, or are being evaluated for, COVID-19. Put on a facemask before you enter the facility. These steps will help the healthcare provider’s office to keep other people in the office or waiting room from getting infected or exposed.

Ask your healthcare provider to call the local or state health department to discuss your situation.

If you have a medical emergency and need to call 911, notify the dispatch personnel that you have, or may have COVID-19. If possible, put on a facemask before emergency medical services arrive.

Discontinuing home isolation

For individuals with symptoms who are confirmed or suspected cases of COVID-19 and are directed to care for themselves at home, discontinue home isolation under the following conditions:

- At least 3 days (72 hours) have passed since recovery defined as resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms (e.g., cough, shortness of breath); AND,
- At least 7 days have passed since symptoms first appeared. Individuals with laboratory-confirmed COVID-19 who have not had any symptoms may discontinue home isolation when at least 7 days have passed since the date of their first positive COVID-19 diagnostic test and have had no subsequent illness.

Additional information for your household members, intimate partners, and caregivers is available at:

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-prevent-spread.html>

Return to Work Criteria for Employees with Confirmed or Suspected COVID-19

Decisions about return to work for employees with confirmed or suspected COVID-19 should be made in the context of local circumstances. Options include a test-based strategy or a non-test-based strategy (i.e., time-since-illness-onset and time-since-recovery strategy).

Use one of the below strategies to determine when employees may return to work in healthcare settings:

Test-based strategy. Exclude from work until resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms (e.g., cough, shortness of breath), and negative results of an FDA Emergency Use Authorized molecular assay for COVID-19 from at least two consecutive nasopharyngeal swab specimens collected ≥ 24 hours apart (total of two negative specimens).

Non-test-based strategy. Exclude from work until at least 3 days (72 hours) have passed since recovery defined as resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms (e.g., cough, shortness of breath); and at least 7 days have passed since symptoms first appeared if employees were never tested for COVID-19 but have an alternate diagnosis (e.g., tested positive for influenza), criteria for return to work should be based on that diagnosis.

**Guidelines are based off the CDC’s interim recommendations. <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html>*